Stress - another possible mechanism of neurotoxicity in neurodegenerative diseases

¹Maria Filomena Xavier Mendes; ² Paulo de Oliveira; ³Maria Solange Gosik; ⁴Renata Garcia Lino; ⁵Raquel Bruno Kalile; ⁶Danielle da Silva barbas; ⁷ Romeu Carillo Jr

- 1 Master in Neurology UFRJ; Coordinator of ABRAH courses in Rio de Janeiro; Professor of ABRAH courses; Invited professor of optional disciplines of homeopathy at UFF; Homeopathic doctor of SMS-RJ-filoabrah@yahoo.com.br
- 2 Professor University of Évora, Portugal; graduated in homeopathy at IMT; Internship at ABRAH in partnership with IMT
- 3 Homeopath and Pediatrician;. National coordinator of ABRAH courses; President of ABRAH; Project Coordination Assistence "cute and healthy" by the Associação Abrigo Rainha da Paz; Homeopath responsible for care in the CAPS for juvenile children Taboão da Serra, SP
- 4 Master in Collective Health UFF; Pediatric and Homeopathy phisician; Supervisor of homeopathy ambulatory of Abrah in Rio de Janeiro; Doctoral student in public health UFRJ.
- 5- Master in in Family Health: Homeopath by ABRAH: Supervisor of homeopathy ambulatory of Abrah in Rio de Janeiro
- 6 Professor University of Brasilia UNB; Coordinator of the pediatric medical residency -HUB; Master in Collective Health-

UFRJ; Doctoral student in public health- UNB; Pediatric and Homeopathy phisician - ABRAH; Professor of ABRAH courses

7 - Head of the Homeopathy Clinic at Hospital do Servidor Público Municipal de São Paulo - HSPM; Course Advisor extension da Abrah - Brazil

Background: The literature associates oxidative stress with the production of free radicals. which lead to neurodegeneration. They present innumerable hypotheses, among which are abnormalities in the functioning of the hypothalamic-pituitary-adrenal axis, neurotoxic effects and neuronal oxidative damage. Clinical observation has shown that in neurodegenerative diseases such as Multiple Sclerosis (MS) and Amyotrophic Lateral Sclerosis (ALS) there is a report of prolonged or violent emotional stress preceding the symptoms. Aims: Using the Carillo Complex Systems Model, present some possibilities on how stress can contribute to neurodegeneration. Methodology: Nine cases of ALS and six cases of MS were evaluated, pathologies already classified as belonging to syphilinism. Literature review on stress and neurotoxicity carried out. Results and discussion: Syphilinism is an instability with a predominantly intrisic origin to the system with a chronic caracter. This diathesis characterized by a dissipative deficiency, predominantly hepatic, to the processing of certain elements or potentially toxic substances with exogenous origin or endogenous Such non-processed substances are unstable factors in the system, with greater affinity for certain tissues, like the nervous system. Among the toxins, we find alcohol, esters, formaldehyde, aloe, ketones, aldehydes, etc. The final hepatic metabolism of cortisol results in cortic acids and cortol, which use the same enzymatic system as alcohol, and can be considered syphilinic toxins. Ethanol can act directly at the circadian rhythm, disrupting it and generating stressful substances such as cortisol, regardless of an external event, increasing the toxin level. The inflammatory process generated by the production of free radicals and metabolic abnormalities, including the reduction of neuropeptide Y that modulates inflammatory activity in the nervous system, leads to changes that can result in neurodegeneration. Conclusion: Inflammation caused by toxins from prolonged/violent emotional stress can lead to several changes in syphilinic individuals, due to failure in the dissipative process, including neurodegeneration.

Keywords: Stress, neurotoxicity, Complex Sistems Model, Syphilinism

References

Carillo Junior, Romeu. Fundamentos da Homeopatia Constitucional: Homeolivros.

Carillo Junior, Romeu. O Milagre da Imperfeição. São Paulo: Cultrix, 2008.

Carillo Junior, Romeu. Homeopatia Medicina interna e Terapêutica. 2° ed. Homeolivros, 2007